



Flector® Tissueal 140 mg Medicated Plaster



Diclofenac sodium (as diclofenac epolamine)
10 medicated plasters

Each dressing is impregnated with gel containing diclofenac epolamine corresponding to 140 mg of diclofenac sodium (1% w/w).
Each 140 cm² (10 cm x 14 cm) medicated plaster contains a total of 180 mg diclofenac epolamine corresponding to 140 mg of diclofenac sodium (E218), propyl parahydroxybenzoate (E216) and propylene glycol

Excipients: methyl parahydroxybenzoate (E218), propyl parahydroxybenzoate (E216) and propylene glycol
Other excipients:
Supporting layer: Unwoven polyester support.
Adhesive layer (active Gel): Gelatin, povidone (K90), liquid sorbitol (non crystallising), heavy kaolin, titanium dioxide (E171), disodium edetate (E385), tartaric acid, aluminum glycidate, carmellose sodium, sodium polyacrylate, 1,3-butylene glycol, polysorbate 80, Dalin PH perfume (propylene glycol, benzylalicylate, phenylethyl alcohol, alpha amyrcinnamic aldehyde, hydroxycyclohexyl, phenylethyl phenylacetate, cinnamyl acetate, benzyl acetate, terpineol, dynamic alcohol), cyclamenaldehyde, purified water.

See leaflet for further information.
Cutaneous use
Keep out of the reach and sight of children.
Do not store above 25°C.
After first opening of the sealed envelope: the product can be stored 3 months maximum.
Remaining active ingredients of the plaster may pose a risk to the aquatic environment. Do not flush used plasters down the toilet. The plaster should be disposed of according to local requirements.

INSTRUCTIONS ON USE

Read the package leaflet before use.
This medicine is indicated for adults and adolescents (aged 16 years or older) for the local symptomatic treatment of pain in epicondylitis (tennis elbow and ankle sprains).
- For elbow pain (epicondylitis), one medicated plaster should be applied in the morning. This should be removed and a fresh plaster applied in the evening. This procedure may be repeated up to a maximum of 14 days duration.
- For ankle sprains one plaster should be used per day up to a maximum of 3 days duration.
- Ankle sprain will normally improve within 3 days and elbow pain within 14 days. If they do not, or if your symptoms get worse, tell your doctor. For adolescents use, see leaflet.
- Do not use it at the same time as other medicines that contain diclofenac or other non-steroidal anti-inflammatory agents, whether these are to be applied to the skin or taken by some other route.
- Do not use the medicated plaster if you are allergic to diclofenac, acetylsalicylic acid or any other NSAID, if you are entering or are already beyond your 6th month of pregnancy, if you currently have a stomach ulcer (peptic ulcer) or if you have any damaged skin including weeping or infected areas, eczema, burns or wounds at the body site where the plaster should be applied, if you are a child or adolescent aged less than 16 years.
- Do not apply the medicated plaster to the eyes, nose, mouth, genital or anal areas. If it does come into contact with these areas, rinse away with water.
- When wearing the plaster, you should avoid exposing it to direct or solarium sunlight in order to reduce the risk of photosensitivity.

LOT : 6754321
EXP : 31.07.2021
04/9/18



Read the package leaflet before use.
This medicine is indicated for adults and adolescents (aged 16 years or older) for the local symptomatic treatment of pain in epicondylitis (tennis elbow and ankle sprains).
- For elbow pain (epicondylitis), one medicated plaster should be applied in the morning. This should be removed and a fresh plaster applied in the evening. This procedure may be repeated up to a maximum of 14 days duration.
- For ankle sprains one plaster should be used per day up to a maximum of 3 days duration.
- Ankle sprain will normally improve within 3 days and elbow pain within 14 days. If they do not, or if your symptoms get worse, tell your doctor. For adolescents use, see leaflet.
- Do not use it at the same time as other medicines that contain diclofenac or other non-steroidal anti-inflammatory agents, whether these are to be applied to the skin or taken by some other route.
- Do not use the medicated plaster if you are allergic to diclofenac, acetylsalicylic acid or any other NSAID, if you are entering or are already beyond your 6th month of pregnancy, if you currently have a stomach ulcer (peptic ulcer) or if you have any damaged skin including weeping or infected areas, eczema, burns or wounds at the body site where the plaster should be applied, if you are a child or adolescent aged less than 16 years.
- Do not apply the medicated plaster to the eyes, nose, mouth, genital or anal areas. If it does come into contact with these areas, rinse away with water.
- When wearing the plaster, you should avoid exposing it to direct or solarium sunlight in order to reduce the risk of photosensitivity.

MA Holder:
IBSA Farmaceutici Italia srl, via Martiri di Cefalonia, 2 - 26900 Lodi, Italy
PL 21039/0004

(UK) 1227



Flector® Tissueal 140 mg Medicated Plaster